



Al-Anbar Medical Journal

Official Journal of Medical College / University of Anbar

Anb Med J

Vol.10 No.1 March 2012



ISSN: 2070 - 8882



Al-Anbar Medical Journal is a Periodic Peer-Reviewed Journal
Published Biannually by the Anbar Medical College.

Editorial Team

Ass. Prof. Salah N. Al-Anli

Ass. Prof. Fakhri Jamil AL-Dalla Ali

Ass. Prof. Ziad H. Abd

General Supervision

Editor In Chief

Executive Editor

Editorial Board

Ass. Prof. Adnan H. Mahdi

Ass. Prof. Majid Abdul Karim Lafi

Ass. Prof. Kahlid Abdullah Al-Rawi

Ass. Prof. Mahasin A. Altaha

Ass. Prof. Mushtaq Talib Salih

Ass. Prof. Tahrir N. Aldelaimi

Dr. Rijaa J. Khalaf

Dr. Arkan A. Jasem A-Iesawi

Dr. Latif A. Abood

Advisory Board

Prof. Jamal Al- Rawl	ph. D comm. Med.
Prof. Rijaa Haql Ismaael	FRCS Faciomaxillary
Prof. Shehab A. Lafi	Ph. D Med. Microbial.
Prof. Ayad Ali Al-Beer	FRCS Urology
Prof. Mahjob Al-Nidawi	MRCP
Prof. Hamed Al-Janabi	MRCP
Ass. Prof. Neama Hamad Al-Heti	FIBMS Urology

Editorial Office

Copy Editor : Bilal Majed Al-Anli

Sponsored By



Dar Al Dawa Scientific Bureau



Contents

Volume 10 , Number 1 , March 2012

Name	No.
Common Rota Virus Gastroenteritis in Children under 5 Years in Maternity and Children Teaching Hospital, western Iraq Qaiss A. Alani, Salih A. Al-Rawi, Ayad K. Salih, Saleem O.G. Al-Mawla	5-11
Management of Pediatric Facial Trauma by Pyriform Suspension Tahrir N. N. Aldelaimi	13-17
Honey depots phagocytosis in vitro Shehab Ahmed Lafi , Huda R. Sabar Al-Dulaimy , Muntaha M. Al-Aloosi.	19-23
A Spectrum Estimation and Clinical Evaluation of Low Back Pain in General Practice Hammodi F J Al-Jumaily	25-35
Cranioplasty: Comparison Between Utilizing Titanium Mesh, Prefabricated Polymethyl methacrylate and Autograft Bone Hisham Maddah Al-Alousi, Kamal T. AL-Kubaisi, Baha'a M. Al-Aani	37-43
Dental Anxiety and it's Relation to Serum Cortisol Level Before Dental Surgical Treatment Dunia W. Alfayad , Eman Majed Rasheed Al-Hadithy	45-50
Kala-azar in Al-Anbar Governorate, Western Iraq Zaid R Al-Ani, Adnan MH Al-Hamwandi, Afrah AS Al-Ma'eeni, Mehdi Kh Al-Ta'aie	52-60
Outcome of Surgical intervention in vascular injuries in Al-Fallujah General Hospital \ 2010 Mohammed Tafash Dagash, Abdulwahhab Abdulrazzaq Mohammed , Yaseen Taha Sarhan	62-70
Inhibition of Uropathogenic Citrobacter Adhesion by Biosurfactants Extracted from Vaginal Lactobacillus acidophilus Omar Abd Ulkareem Ali	72-80
Prevalence of Oesophago - Gastroduodenal Illnesses Among Dyspeptic Children in Al-Anbar Governorate Abdulla Eltaif Jassim	82-88
A Comparison of Arch Width in Adults with Normal Class I Occlusion and Adults with Class II Division I Malocclusion in Ramadi City Zena Hekmat Al-Taei	90-95
Case Report: Scrotal Gigantism Niema H. Al-Heeti, Abdul-Fatah N. Al-Ani	97-101



Dental Anxiety and it's Relation to Serum Cortisol Level Before Dental Surgical Treatment

Dunia W. Alfayad¹, Eman Majed Rasheed Al-Hadithy²

Lecturer oral pathology Department 1; Assistant lecturer Basic Sciences Department 2 ; Faculty of Dentistry; University of Anbar, Iraq.

Abstract

Background: Dental anxiety is still prevalent, despite advances in treatment methods, and it affects the utilization of health does not only decrease the pain threshold, but actually leads to the perception of painless stimuli as painful. services. Cortisol is an important hormone in the body, secreted by the adrenal glands and involved in many functions.

Objectives: To evaluate the relation between the level of serum cortisol and increase anxiety in dental patients before minor oral surgery.

Materials and Method: Sixty patients were included in this study were attended Alkatana Specialized Dental Center from October/4/2010 till December/1/2010, their age ranged from (16-54 years) 27 patients were females and 33 were males. They were divided into two groups, 30 of them as controls (they didn't need any dental surgery) who match the other 30 patients study group in age, sex and their general health status but they needed minor oral surgery. Blood samples were collected from all patients between 10-11 Am., and about 5 minutes before surgery to the patients of the study group. Serum cortisol level was measured by using radioimmunoassay analysis.

Results: A total of 60 patients were enrolled in this study. Regarding the control group 15 were females and 15 were males while for the study group patients 12 were females and 18 were males. Serum cortisol level was significantly different between two groups the mean was 13.05 ± 6.51 for control patients and 23.62 ± 10.12 respectively and the Coefficient correlation (r) between serum cortisol level and pulse rate in both groups were 0.16 ($p > 0.05$) for the control patients and 0.58 ($p < 0.01$) for the study group patients. When serum cortisol concentrations in study group were distributed according to the age of the sample, there was a highly significant positive correlation between these variables ($r = 0.36$, $p < 0.05$). Also pulse rate in study group was found highly positive association with age ($r = 0.55$, $p < 0.01$).

Conclusions: It is concluded that the study group patients exhibited significantly higher levels of serum cortisol and pulse rate than that of the control group. This deference is suggested to be due to phobia from dental surgical work. As recommendation we suggest giving patient 5 mg of diazepam at the night before the operation.

Keywords: dental anxiety; pain; cortisol.

Corresponding author: Dunia W. Alfayad, E-Mail: DuniaAl-Fayad@yahoo.com, Tel. 07800700757

Introduction:

Dental treatment may cause pain and discomfort. Even the expectation of pain increases dental anxiety, thus maintaining the number of dentally anxious persons ^(1,2).

Dental fear, anxiety, and phobia have consistently been reported as widespread problems that persist despite the technological advances that have made dentistry less painful and less uncomfortable.



It is well documented that dental fear has a significant impact on dental care utilization behaviors^(3,4,5).

Anxiety not only lowers the pain threshold, but actually leads to the perception of painless stimuli as painful. Cortisol is an important hormone in the body, secreted by the adrenal glands and involved in many functions like: Proper glucose metabolism; Regulation of blood pressure; Insulin release for blood sugar maintenance; Immune function and inflammatory response⁽⁶⁾.

Prolonged high levels of cortisol can lead to heart disease and other health problems. Treatments for dental fear often include a combination of behavioral and pharmacological techniques⁽⁷⁾.

The Aim of the Study is To evaluate the level of serum cortisol in dental patients before minor oral surgical procedure and it is relation to increase anxiety.

Materials and Method:

Sixty patients were included in this study were attended Alkatana Specialized Dental Center from October/4/2010 till December/1/2010, their age ranged from (16-54 years) 27 patients were females and 33 were males. They were divided into two groups, 30 of them as controls (they didn't need any dental surgery) who match the other 30 patients study group in age, sex and their general health status but they needed minor oral surgery. Single female experimenter, measured blood pressure and pulse rate for all the patients and then she started to collect blood samples from them between 10-11 Am., about 5 minutes before surgery to the patients of the study group.

A 5 ml. blood samples were placed immediately in heparinized tubes containing sodium meta bisulfate as an antioxidant then centrifuged at 5000 r.p.m. for three minutes.

Plasma samples were immediately pipette into tubes containing 0.5 ml. of iced 4N per chloric acid and then frozen. Serum cortisol level was measured by radioimmunoassay analysis using Cortisol Ria Kit / Iso date 20 / 20 Gamma Center / USA according to manufacture instructions.

For Statistical analysis, Chi-square tests (χ^2) with Yates' correction were used to compare the distribution of subjects in the categorical variables. Data of serum cortisol levels are presented as mean values and standard deviation (SD). Serum cortisol concentrations were log-transformed, which normalized distribution of cortisol measures. Student's *t*-tests were used to compare continuous variables. Correlations were explored by means of the Pearson correlation coefficient. Correlations involving pulse rate and serum cortisol level were analyzed by means of Spearman correlation coefficient. Statistical significance was accepted at $P < 0.05$.

Results:

A total of 60 patients were enrolled in this study. Regarding the control group 15 were females and 15 were males while for the study group patients 12 were females and 18 were males. Serum cortisol level was significantly different between two groups the mean was 13.05 ± 6.51 for control patients and 23.62 ± 10.12 respectively and the Coefficient correlation (*r*) between serum cortisol level and pulse rate in both groups were 0.16 ($p > 0.05$) for the control patients and 0.58 ($p < 0.01$) for the study group patients as shown in figures 1 and 2.

When serum cortisol concentrations in study group were distributed according to the age of the sample, there was a highly significant positive correlation between these variables ($r = 0.36$, $p < 0.05$). Also pulse rate in study group was found highly positive association with age ($r = 0.55$, $p < 0.01$).




Figure-1 shows the relation between pulse rate and serum cortisol level in control group patients.



Figure-2 shows the relation between pulse rate and serum cortisol level in study group patients.



Discussion:

In present study show dental anxiety or stress increases serum cortisol secretion as a result of increased corticotrophin releasing hormone (CRH) secretion and increase sympathetic tone therefore, cortisol play a role in body response to stress (activation of energy metabolism, increase in cardiac performance). Also cortisol increase myocardial contractility and vasoconstriction, these are described as permissive effects of cortisol. Cortisol increases the synthesis of epinephrine in the medulla & angiotensinogen in the liver⁽⁸⁾. Researchers such as Chapman and Kirby-Turner (1999) suggest that there is a relationship between general and dental anxiety⁽⁹⁾. It should be borne in mind that the origins of dental anxiety are numerous and complex and have been shown in other parts of the world to be associated with a, irregular attendance pattern, history of extractions, having a dentally anxious parent and is also related to memory distortions and personality types. Highly anxious dental patients tend to overestimate the intensity of aversive dental events even if they have never experienced the particular experience before^(10,11,12,13). Emergency dental situations involve a high prevalence of fear and anxiety, and it is important for the dentist to be able to recognize such patients⁽¹⁴⁾. Normally, cortisol present in the body at higher levels in the morning, and at its lowest at night. Although stress isn't the only reason that cortisol is secreted into the blood stream, it has been termed "the stress hormone" because it's also secreted in higher levels during the response to stress, and is responsible for several stress-related changes in the body. Cortisol potentiates production of Catecholamines and regulates beta-adrenergic receptors^(15,16,17).

Direct experience is the most common way make people develop dental fears. Dental anxiety has been associated with avoidance of dental care and affects not only general health, but also it is lead to the sleep disturbance, and social interactions with work performance⁽¹⁸⁾. People who are dentally fearful tend to avoid treatment and are much more likely to attend the dentist when prompted by pain and their avoidance removes opportunities for learning, as does the dental profession's emphasis on the use of sedation or general anesthetic with fearful patients^(19,20,21). Milgrom et al. reported that younger subjects were more anxious than older ones⁽²²⁾, while, in the present study dental anxiety was correlated with age. This may be due to small number of patient's age more than 50 who seek care. Zimmer et al. and Kirsch Baum et al. demonstrated that the pain is a stimulus that results in significant increases of salivary cortisol⁽²³⁾. The present results are in concordance with these studies, in which dental anxiety in study group patients exhibited significantly higher levels of serum cortisol due to phobia from dental surgical work because, stress increases serum cortisol secretion and increase sympathetic tone.

Moreover, the pulse rate in study group was found highly positive association with age immediately prior to the dental checkup. This is in agreement with Rayen et al. the results clearly showed a significant change in the systolic pressure and heart rate in all the situations in the dental operatory area⁽²⁴⁾. However the present study showed a significant increase in pulse rate in almost all the dental procedures when compared with normal. It could be because the treatment had been started without any behavior management.



Conclusions and Recommendations:

It is concluded that the study group patients exhibited significantly higher levels of serum cortisol and pulse rate than that of the control group. This deference is suggested to be due to phobia from dental surgical work. As recommendation we suggest giving patient 5 mg of diazepam at the night before the operation.

References

1. Locker D. (2003) psychosocial consequences of dental fear and anxiety. *Community Dent Oral Epidemiol*; 31: 144-51.
2. Hagglin C, Hakeberg M, Ahlqvist M, Sullivan M, Berggren U. (2000) Factors associated with dental anxiety and attendance in middle aged and elderly women. *Community Dent Oral Epidemiol*; 28:451-60.
3. Skaret E, Raadal M Kvale G, Berg E. (2000) Factors related to missed and cancelled dental appointments among adolescents in Norway. *Eur J Oral Sci*; 108:175-83.
4. Good BJ, Kleinman AM (1985) Culture and anxiety: cross-cultural evidence for the patterning of anxiety disorders. In *Anxiety and the Anxiety Disorders*. Edited by: Tuma AH, Maser J. Hillsdale NJ: Lawrence Earlbaum;297-323.
5. Bracha HS, Vega EM, Vega CB (2006). "Posttraumatic dental-care anxiety (PTDA): Is "dental phobia" a misnomer?" *Hawaii Dent J* 37 (5): 17-9.
6. Pitman, R.K. & Delahanty, D.L. (2005). Conceptually driven pharmacologic approaches to acute trauma. *CNS spectr*, 10, 99-106.
7. American Academy of Pediatric Dentistry (AAPD) (2006). Guideline on behavior guidance for the pediatric dental patient.
8. Color Atlas of physiology (2003).
9. Chapman, HR & Kirby-Turner.(1999) NC Dental fear in children—a proposed model *British Dental Journal*; 187(8): 408-12
10. Dixon GS, Thomson WM, Krueger E. (2003) The west coast study. Itself reported dental Skaret E,Raadal M,Kvale G,Berg E.Gender based differences in factors related to non-utilization of dental care in young Norwegians. A longitudinal study.*Eur J Oral Sci*;111: 377-82.
11. Steele JG,WallsAW, AyatollahiSM, Murray JJ. (1996) Dental attitudes and behavior among a Sample Of dentate older adults from three English Communities.*Br Dent J*;180:131-6.
12. Kvale, G et. al.(2004) Dental fear in adults: a meta-analysis of behavioral interventions *Community Dentistry and Oral Epidemiology*; 32: 250-264
13. Mac Entee MI, Stolar E, Glick N. (1993) Influence of age and gender on oral health and related behavior in an independent elderly population. *Community Dent Oral Epidemiol*;21:234-9.
14. Erten H,Akarslan ZZ, Bodrumlu E. (2006) "Dental fear and anxiety levels of patients attending a dental clinic". *Quintessence Int* 37 (4): 304-10.
15. King SL, Hegadoren KM. (2002) Stress hormones: how do they measure up? *BiolRes Nurs* 4, 92-103.
16. Manor I, Tyano S, Mel E, et al. (2002). "Family-based and association studies of monoamine oxidase A and attention deficit hyperactivity disorder (ADHD): preferential transmission of the long promoter-region repeat and its association with impaired performance on a continuous performance test (TOVA)". *Molecular Psychiatry* 7 (6): 626-32.
17. Brunner HG. (1996) "MAOA deficiency and abnormal behavior: perspectives on an association". *Ciba Foundation Symposium* 194: 155-64; discussion 164-7.
18. Cohen SM, Fiske J,Newton JT (2000) The impact of dental anxiety on daily living.*Br Dent J* 189,385-390.
19. Manski RJ, Moeller JF, Maas WR.(2001) Dental services — an analysis of utilization over 20 years. *J Am Dent Assoc* -132:655-64.
20. Manski RJ, Gold Farb MM.(1996) Dental utilization for older Americans aged 55 - 75.*Gerodontology*;13:49-55.
21. Mac Entee MI, Stolar E, Glick N. (1993) Influence of age and gender on oral health and related behavior in an independent elderly population. *Community Dent Oral Epidemiol*; 21: 234-9.
22. Milgrom P, Fiset L, Melnick S, Weinstein P. (1988) the prevalence and practice management consequences of dental fear in a major US city. *J Am Dent Assoc* 116, 641-647.



23. Zimmer C, Basler HD, Vedder H, Lautenbacher S. (2003) Sex differences in cortisol response to noxious stress. *Clin J Pain* 49, 233-239.
24. Rayen R, Muthu MS, Rao CR, Sivakumar N. Evaluation of physiological and behavioral measures in relation to dental anxiety during sequential dental visits in children. *Indian J Dent Res [serial online]* 2006 [cited 2011 Dec 27];17:27-34.

المحتويات

المجلد ١٠ ، العدد ١ ، آذار ٢٠١٢

الرقم	اسم البحث
١١-٥	دور فايروس الروتا كمسبب للإسهال لدى الأطفال تحت سن الخامسة من العمر في مستشفى النسائية و الأطفال التعليمي في الرمادي قيس عليوي العلي . صالح الراوي . لياد خلف صالح. سليم المولى
١٧-١٣	المعالجة الجراحية لكسور عظام الوجه عند الأطفال تحرير نزال الدليمي
٢٣-١٩	العسل ينشط البلعمة في المختبر شهاب احمد لافي . هدى رافع صبار الدليمي . منتهى مداح الالوسي
٣٥-٢٥	تقدير طيفي و تقييم سريري لأنسج اسفل الظهر لدى العموم حمودي فرحان جاسم الجميلي
٤٣-٣٧	تقويم القحف: دراسة مقارنة بين استخدام شبكة التيتانيوم ومادة البولي مثيل ميثا اكريليت المعدة مسبقا والعظم ذاتي الطعم هشام مداح الالوسي ، كمال تركي الكبيسي ، بهاء مشرف عبد السلام العلي
٥٠-٤٥	القلق النفسي لدى مرضى الاسنان وعلاقة هرمون الكورتيزول قبل عمليات الاسنان الصغرى دنيا وديع الفياض ، ايمان ماجد رشيد
٦٠-٥٢	الحمى السوداء في محافظة الانبار ، غرب العراق زيد رشيد العاني ، عدنان محمد حسن ، افراح عبد المهدي صالح ، مهدي خلف الطائي
٧٠-٦٢	نتائج التداخل الجراحي في إصابات الأوعية الدموية في مستشفى الفلوجة العام / ٢٠١٠ محمد طفش دغش ، عبد الوهاب عبد الرزاق ، ياسين طه سرحان
٨٠-٧٢	تنشيط التصاق بكتريا Citrobacter المسببة للالتهابات المجاري البولية باستخدام محلول Biosurfactants المستخلص من Lactobacillus acidophilus عمر عبد الكريم
٨٨-٨٢	مدى انتشار امراض المريء والمعدة والاثني عشري بين الاطفال المصابين بسوء الهضم في محافظة الانبار ، غرب العراق عبد الله الطيف جاسم
٩٥-٩٠	مقارنة عرض القوس السني للبالغين من ذوي الإطباق من الصنف الأول مع ذوي سوء الإطباق من الصنف الثاني . النموذج الأول في مدينة الرمادي زينة حكمت بشير
١٠١-٩٧	كيس الصفن العملاق - تسجيل حالة نعمة حمد الهيتي ، عبد الفتاح العاني

مجلة الانبار الطبية هي مجلة دورية محكمة تصدر عن كلية الطب
جامعة الانبار وتنتشر أعدادها بمعدل مرتين في السنة.

فريق التحرير

أ.م.د. زياد حماد عبد

أ.م.د. فخري جميل الدلة علي

أ.م.د. صلاح نوري احمد العاني

المدير التنفيذي

رئيس التحرير

الإشراف العام

هيئة التحرير

أ.م.د. خالد عبد الله نافع

أ.م.د. مجيد عبد الكريم لافي

أ.م.د. عدنان حماد مهدي

أ.م.د. تحرير نزال نايف

أ.م.د. مشتاق طالب صالح

أ.م.د. محاسن علي عبد الجبار

د. لطيف احمد عبود

م.د. اركان عبيد جاسم

م.د. رجاء جهاد خلف

الهيئة الاستشارية

أ.د. جمال الراوي

أ.د. محبوب النداوي

أ.د. رجاء حقي اسماعيل

أ.د. اياد علي البير

أ.د. شهاب احمد لافي

أ.د. حامد الجنابي

أ.م.د. نعمه حمد الهيبي

سكرتير هيئة التحرير

بلال ماجد طه العاني

الراعي الرسمي



مكتب دار الدواء العلمي





مجلة الأنبار الطبية

نصدر عن كلية الطب / جامعة الأنبار

Anb Med J

المجلد العاشر - العدد الأول - آذار 2012



ISSN: 2070 - 8882